

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

BT

(19)



Eur päisches Pat ntamt

Europ an Patent Office

Offic uropé n des br vets



(11)

EP 0 882 408 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
09.12.1998 Bulletin 1998/50

(51) Int Cl.⁶: **A23L 1/236**

(21) Application number: **98304315.9**

(22) Date of filing: **01.06.1998**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **02.06.1997 JP 157321/97**
07.11.1997 JP 320519/97

(71) Applicant: **KABUSHIKI KAISHA HAYASHIBARA
SEIBUTSU KAGAKU KENKYUJO**
Okayama-shi Okayama (JP)

(72) Inventors:
• **Chaen, Hiroto**
Okayama-shi, Okayama (JP)
• **Mitsuhashi, Masakazu**
Okayama-shi, Okayama (JP)
• **Miyake, Toshio**
Okayama-shi, Okayama (JP)

(74) Representative: **Daniels, Jeffrey Nicholas et al**
Page White & Farrer
54 Doughty Street
London WC1N 2LS (GB)

(54) **Method for improving the aftertaste of sucrose, and uses thereof**

(57) A method for improving the aftertaste of sucrose without lowering the sweetening power of sucrose, which comprises incorporating trehalose into sucrose in an amount of 2-12% to the sucrose, on a dry

solid basis. The method produces saccharide compositions and food products having improved aftertaste of sucrose without lowering the sweetening power of sucrose.

EP 0 882 408 A1

Description

The present invention relates to a method for improving the aftertaste of sucrose and uses thereof, and more particularly to a method for improving the aftertaste of sucrose characterized in that it comprises incorporating trehalose into sucrose in an amount of 2-12 w/w % to the sucrose, on a dry solid basis (d.s.b.), and saccharide compositions and food products, wherein the aftertaste of sucrose is improved by using the method without lowering the sweetening power of sucrose.

Sucrose, a natural sweetener used for a long time, has become to be noticed of its thick taste, as a main drawback, which causes unpleasant long-lasting aftertaste, in modern diet styles flooded with food materials and fast foods.

To improve the drawback, for example, Japanese Patent Kokoku No. 44,476/86 proposed a method for improving the aftertaste of sucrose by treating sucrose solutions with ultrasonic; the method is scarcely used because of its relatively-low effect for its complicatedness. The combination use of sucrose with other saccharides such as maltose, trehalose, and maltotetraose is known in the food industries as disclosed, for example, in Japanese Patent Kokai No. 319,486/94 and Japanese Patent Kokoku Nos. 77,381/93 and 89,916/95. These other saccharides are low-sweetened sweeteners with a reduced sweetening power of sucrose. On the pretext of fitting to modern feelings, these conventional methods aim to lower the content of sucrose in food products in order to reduce the sweetening power as much as possible and increase the content of total saccharides as high as possible within the margin of the reduced sweetening power, resulting in a relatively-long shelf-life of the food products. Usually, the other saccharides are added to sucrose in an amount of about 40-300 w/w % to the sucrose, d.s.b. In some cases, such use of sucrose hinders the effective high-sweetening power of sucrose, and in many cases, it deteriorates the flavor and taste of sucrose. It has been strongly demanded to improve the aftertaste of sucrose as a drawback while retaining the sweetening powder, flavor, and taste.

The object of the present invention is to provide a method for improving the aftertaste of sucrose due to its thick taste without substantially lowering the sweetening power of sucrose, and uses thereof.

To attain the above objects, the present inventors energetically studied for the combination use of sucrose and other saccharides. As a result, they unexpectedly found that the present object is attained by incorporating trehalose, α,α -trehalose, into sucrose in a specific proportion, and accomplished this invention; the inventors found that incorporation of trehalose into sucrose in an amount of 2-12 w/w % (hereinafter the wording "w/w %" is abbreviated as "%" throughout the specification, unless specified otherwise) to the sucrose, d.s.b., eliminates the thick taste and aftertaste of sucrose without substantially lowering the sweetening power of sucrose, and they accomplished this invention which is mainly constructed by:

- (1) A method for improving the aftertaste of sucrose, characterized by incorporating trehalose into sucrose in an amount of 2-12% to the sucrose, d.s.b.;
- (2) A saccharide composition which contains trehalose in an amount of 2-12% to sucrose, d.s.b.;
- (3) A food product with improved aftertaste of sucrose without lowering the sweetening power of sucrose, where trehalose is incorporated into sucrose in an amount of 2-12% of the sucrose, d.s.b., without substantially lowering the sweetening power of sucrose; and
- (4) An agent for improving the aftertaste of sucrose, which comprises trehalose as an effective ingredient.

The present invention will now be described by way of example only.

Any types and forms of sucrose can be used in the present invention independently of their origins as long as the present invention improves their taste quality, particularly, their aftertaste. For example, white sugars, granulated sugars, white superior soft sugars, powdered sugars, frosted sugars, and liquid sugars can be used. Saccharides comprising sucrose and other saccharides can be also used. In general, those rich in sucrose can exert more easily the effect of the present invention, and the following saccharide compositions can be advantageously used in the present invention; black sugars containing sucrose and small amounts of invert sugars and minerals, α -galactosyl sucrose such as raffinose and stachyose, saccharides containing sucrose, α -glycosyl sucrose such as erlose and maltosylsucrose, saccharides containing maltooligosaccharides and sucrose, i.e., "COUPLING SUGAR®, β -galactosylsucrose, i.e., lactosucrose, and saccharides containing lactose and sucrose, i.e., NYUKAOLIGO®".

Any types and forms of trehalose can be used in the present invention independently of their origins as long as they can improve the flavor and taste of sucrose, particularly, the aftertaste of sucrose. For example, trehalose derived from starches as disclosed in Japanese Patent Kokai Nos. 319,486/94 and 213,283/95, and trehalose from maltose as disclosed in Japanese Patent Kokai Nos. 216,695/83 and 170,977/95 can be advantageously used. The sweetening power of trehalose is known to be an about 45% sweetening power of sucrose and is not changed independently of the form of a crystalline hydrate, crystalline anhydride, powder, and liquid. The form of trehalose can be appropriately chosen as occasion demand. Trehalose can be advantageously used as an effective ingredient for agents for improving the aftertaste of sucrose.

To attain the present object, trehalose should be incorporated into sucrose in an amount of 2-12% to the sucrose, d.s.b. If the content is below 2%, d.s.b., it improves insufficiently the aftertaste of sucrose, while the content is over 12%, d.s.b., it unsatisfactorily reduces the sweetening power of sucrose. As long as not inhibiting the present object, other saccharides excluding sucrose and trehalose can be coexisted: For example, monosaccharides such as glucose, fructose, and galactose; disaccharides such as maltose, isomaltose, lactose, and palatinose; trisaccharides such as maltotriose, erlose, raffinose, lactosucrose, and ketose; tetrasaccharides such as maltotetraose and stachyose; and sugar alcohols such as xylitol, sorbitol, maltitol, lactitol, and maltotriitol, all of which are appropriately chosen, particularly, saccharide compositions containing oligosaccharides having a sucrose structure can be advantageously used. The present saccharide compositions thus obtained contain trehalose in an amount of 2-12% to sucrose, d.s.b. The higher the content of sucrose, the more easily exerted the present effect; Preferably, those containing at least 10%, and more preferably, at least 40% sucrose, d.s.b., can be suitably used independently of their forms of solid, paste, and liquid.

The present saccharide compositions thus obtained have satisfactorily improved aftertaste of sucrose free from the reduction of sweetening power, and they can be arbitrarily and widely used to sweeten food products. Specifically, the saccharide compositions exert their effect in food products which contain relatively-large amounts of the compositions, and preferably they can be advantageously used in food products containing at least 30% of the compositions, d.s.b.: For example, seasonings such as sweeteners, table sugars, coffee sugars, "mentsuyu" (a sauce for Japanese vermicelli), sauces, catsups, "yakniku-no-tare" (a sauce for Japanese grilled meat), instant soup mixes, and complex seasonings; Japanese confectioneries and material saccharides thereof such as *an* (a bean jam), "uiro" (a sweet rice jelly), "yokan" (a sweet jelly of beans), jellies, "kingyoku" (a kind of jelly), *pao de Castellas*, and "amedama" (a Japanese toffee); Western confectioneries and material saccharides thereof such as buns, cookies, pies, puddings, chewing gums, chocolates, sand creams, custard creams, butter creams, cream puffs, wafers, sponge cakes, flour pastes, peanut pastes, candies, spreads, "toka" (conserves), syrups, bonbons, and sweetened condensed milks; soft drinks and material saccharides thereof such as coffees, cocoas, teas, juices, sour milk beverages, and beverages containing lactic acid bacteria; ice candies and material saccharides thereof such as ice creams, sherbets, "kaki-gori", and "korigomitsuru"; instant food products and material saccharides thereof such as instant juices, instant coffees, instant cocoas, instant teas, instant "shiruko", and instant puddings; and other food products such as foods for infants, therapeutic foods, beverages with supplemental nutritions, dried foods, and "chinmi" which are sweetened with the present saccharide compositions.

When using powdered sucrose as sucrose in the process for producing the present saccharide compositions, the addition of powdery trehalose to the powdered sucrose in an amount of 2-12% to the powdered sucrose, d.s.b., improves the aftertaste of sucrose without substantially lowering the sweetening power of sucrose and exerts solidification-preventing effect on the powdered sucrose.

The powdery saccharide compositions which are prevented from solidification can be advantageously used in food products in general and used as material saccharides in chewing gums, chocolates, sand creams, icings, sugar coatings, and toppings, all of which prefer powdery saccharides.

Methods for incorporating the present saccharide compositions in the aforesaid food products are those which can incorporate thereunto the saccharide compositions during the processes before completion of their processings; conventional methods such as mixing, kneading, dissolving, melting, soaking, permeating, sprinkling, applying, coating, spraying, injecting, crystallizing, and solidifying can be appropriately chosen. If necessary, other materials including appropriate materials and additives such as different sweeteners, taste-imparting seasonings, flavoring agents, coloring agents, emulsifiers, nutrients, antioxidants, and fillers can be arbitrarily incorporated into food products in adequate amounts.

The following experiments describe the present invention:

Experiment 1

Influence of saccharides on both the sweetening power and the improvement of the aftertaste of sucrose

The influence of coexisting saccharides on both the sweetening power and the improvement of the aftertaste of sucrose was studied. The saccharides used were commercially available ones in reagent grade. A 10% aqueous sucrose solution was used as a control solution, and a 10% aqueous solution containing 10% each of the saccharides, d.s.b., was used as a test solution. The saccharides were glucose, sorbitol, maltose, maltitol, lactose, trehalose or α , α -trehalose, and neotrehalose or α , β -trehalose. A panel test was conducted using the paired difference test which compared the sweetening power and aftertaste of sucrose in the control and test solutions. The panel test was conducted at a room temperature of 25°C using 13 volunteers consisting of eight males and five females. The sweetening power was evaluated by allowing the volunteers to compare the test solution with the control solution and to select their answers among the grades "inferior", "equal" and "superior", while the improvement of the aftertaste of sucrose

EP 0 882 408 A1

was evaluated by allowing the volunteers to compare the test solution with the control solution and to select their answers among the grades "inferior", "equal" and "superior", followed by counting the number of volunteers who selected the grades each. The results are in Table 1:

5

10

15

20

25

30

35

40

45

50

55

Table 1

Saccharide coexisted in sucrose	Grade	Sweetening power	Aftertaste	Judgement
Glucose	C>T	11	1	Control
	C=T	2	10	
	C<T	0	2	
Sorbitol	C>T	11	1	Control
	C=T	2	11	
	C<T	0	1	
Maltose	C>T	13	1	Control
	C=T	0	10	
	C<T	0	2	
Maltitol	C>T	9	2	Control
	C=T	4	9	
	C<T	0	2	
Lactose	C>T	13	1	Control
	C=T	0	11	
	C<T	0	1	
Trehalose (α , α -trehalose)	C>T	2	0	Present invention
	C=T	10	1	
	C<T	1	12	
Neotrehalose (α , β -trehalose)	C>T	13	2	Control
	C=T	0	9	
	C<T	0	2	

Note : In the table, the numerals are the number of volunteers among 13, who selected the grades each.
The symbols "C>T", "C=T", and "C<T" mean that the test solution was inferior to, equal to, and superior to the control solution with only sucrose.

As evident from the results in Table 1, it was revealed that unlike the sucrose solutions coexisted with other saccharides, the solutions, in which trehalose or α,α -trehalose coexisted with sucrose, had satisfactorily-improved after-taste of sucrose or sharpened aftertaste of sucrose. The coexistence of saccharides other than trehalose lowered the sweetening power of sucrose but could not improve the aftertaste of sucrose with respect to the sharpness. The data shows that trehalose improves the sharpness of aftertaste of sucrose without substantially lowering the sweetness of sucrose, and the effect is characteristic of trehalose.

Experiment 2

Influence of the concentration of trehalose on both the sweetening power and the improvement of the aftertaste of sucrose

The influence of the concentration of trehalose on both the sweetening power and the improvement of the aftertaste of sucrose was studied with an aqueous solution, coffee, and tea. Similarly as the method in Experiment 1, commercially available sucrose and trehalose in reagent grade were used.

In the case of an aqueous solution, a 10 w/v % aqueous solution containing a saccharide composition, containing sucrose and 1.0, 2.0, 5.0, 8.0, 12.0, 16.0 or 20.0% trehalose to the sucrose, d.s.b., was used as a test solution. As a control solution, a 10 w/v % sucrose solution with only the saccharide was used. These test and control solutions with ambient temperature were used in a panel test.

In the case of a coffee as a test solution, commercially available regular coffee powder was extracted in a conventional manner, and in the extract was dissolved 10 w/v % either of the saccharide compositions having the same proportions of sucrose and trehalose as used in the aqueous solution. As a control solution, a 10 w/v % sucrose solution with only the saccharide was used. These test and control solutions in hot conditions were used in a panel test.

In the case of a tea as a test solution, commercially available tea bag was extracted in usual manner, and in which was dissolved 5 w/v % either of the saccharide compositions having the same proportions of sucrose and trehalose as used in the aqueous solution. As a control solution, a 5 w/v % sucrose solution with only the saccharide was used. These test and control solutions in hot conditions were used in a panel test.

Similarly as in Experiment 1, a panel test was conducted using the paired difference test for evaluating the sweetening power and the improvement of the aftertaste of sucrose.

The results are in Table 2:

Table 2

Trehalose content (%) to sucrose, on a dry solid basis	Grade	Aqueous solution*		Coffee in hot		Tea in hot		Judgement
		A	B	A	B	A	B	
1.0	C>T	1	2		3	1	4	Control
	C=T	11	8	2	7	10	7	
	C<T	1	3	1	3	2	2	
2.0	C>T	1	1	1	1	1	0	Present invention
	C=T	10	2	11	2	11	2	
	C<T	2	10	1	10	1	11	
5.0	C>T	0	1	0	1	0	0	Present invention
	C=T	11	2	11	1	10	0	
	C<T	2	10	2	11	3	13	
8.0	C>T	2	0	1	0	0	0	Present invention
	C=T	10	1	11	1	13	2	
	C<T	1	12	1	12	0	11	
12.0	C>T	2	0	1	0	2	1	Present invention
	C=T	10	2	12	1	10	2	
	C<T	1	11	0	12	1	10	
16.0	C>T	4	0	4	1	2	2	Control
	C=T	9	1	8	3	9	3	
	C<T	0	12	1	9	2	8	
20.0	C>T	9	0	11	1	4	2	Control
	C=T	4	0	2	4	7	4	
	C<T	0	13	0	8	2	7	

Note : In the table, the numerals are the number of volunteers among 13, who selected the grades each. The symbols ">", "<" and "<=" mean "ambient temperature", "sweetening power" and "aftertaste", respectively. The symbols "C>T", "C=T", and "C<T" mean that the test solution was inferior to, equal to, and superior to the control solution with only sucrose.

As evident from the results in Table 2, it was revealed that the coexistence of trehalose in an amount from 2.0% to 12.0% to sucrose, d.s.b., independently of aqueous solution, coffee, and tea, improves the aftertaste of sucrose, i. e., the sharpness of the aftertaste of sucrose without substantially lowering the sweetening power of sucrose.

Experiment 3

Solidification-preventing activity of trehalose on powdered saccharides

The solidification-preventing activity of powdery trehalose on a powdery saccharide or a powdered sucrose was studied. One hundred parts by weight of the powdered saccharide or the powdered sucrose, which 90% or higher passes through a JIS 250 μ sieve, prepared by pulverizing granulated sugar, and 10 parts by weight of powdery trehalose prepared by heating "TREHAOSE®", a food grade crystalline trehalose hydrate commercialized by Hayashibara Shoji, Inc., Okayama, Japan, to convert it into an anhydrous crystalline form, and pulverizing the anhydride similarly as in the above granulated sugar, were mixed to homogeneity to obtain a saccharide composition containing trehalose in an amount of about 10% to sucrose, d.s.b. Twenty-kilogram aliquots of the saccharide composition were in a conventional manner injected into polyethylene bags which were then fastened their openings with rubber bands and placed on a three-layered craft paper, followed by sewing the edges of the paper by a machine to enclose the bags for test samples. Five bags of the test samples were placed on a pallet in one layer and loaded with a pallet loaded with 50 bags, one ton in total, containing 20 kg food grade trehalose each, followed by storing in a storage for one month at ambient temperature. As a control, only the powdered sugar or the powdered sucrose was injected into bags similarly as above and stored in the storage. After one-month standing, these bags were opened for examining the conditions of the contents, revealing that the powdered saccharide as the control solidified rigidly but the present saccharide composition was kept in a free-flowing powder, which scarcely solidified but easily collapsed only by shaking by hands. Thus, it was revealed that the saccharide compositions according to the invention have improved aftertaste of sucrose without lowering the sweetening power of sucrose and exert solidification-preventing effect.

Examples A and B disclose the preferred embodiments of the saccharide compositions and food products according to the present invention, respectively:

Example A-1

A saccharide composition was obtained by mixing to homogeneity 100 parts by weight of white superior soft sugar with 10 parts by weight of "TREHAOSE®", a food grade crystalline trehalose hydrate commercialized by Hayashibara Shoji, Inc., Okayama, Japan, which was equal to about 9% of the sucrose, d.s.b., with respect to the trehalose content. The product, having improved aftertaste of sucrose without lowering the sweetening power, can be arbitrarily used to sweeten food products.

Example A-2

One hundred parts by weight of granulated sugar was placed in a fluidized-bed granulator which was then set to give a blowing air temperature of 40°C, followed by granulating the contents while being sprayed with 10 parts by weight of 50% aqueous trehalose solution, which was equal to about 5% of the sucrose, d.s.b., with respect to the trehalose content. The granulated contents were transferred to an aging tower and aged at 25°C for eight hours into a saccharide composition. The product, a stable and free-flowing saccharide composition of granulated sugar coated with trehalose, improved the sharpness of aftertaste of sucrose without lowering the sweetening power so that it can be arbitrarily used in food products.

Example A-3

A saccharide composition was obtained by pulverizing a granulated sugar into a powdered sugar, i.e., a powdered sucrose which 90% or more passes through a JIS 250 μ sieve, and mixing to homogeneity 100 parts by weight of the powdered sugar with five parts by weight of a powdered trehalose, which was equal to about 5% of the sucrose with respect to the trehalose content, d.s.b., and obtained by heating "TREHAOSE®", a food grade crystalline trehalose hydrate, to convert it into an anhydrous form, and pulverizing the crystalline trehalose anhydride similarly as the above granulated sugar. Since the product improved the sharpness of aftertaste of sucrose without lowering the sweetening power of sucrose, it can be arbitrarily used to sweeten food products and suitably used in chewing gums, chocolates, and sand creams as a material saccharide, a powdered saccharide free of solidification and having satisfactory free-flowing ability.

Example A-4

A liquid saccharide composition was obtained by mixing and dissolving one part by weight of "TREHAOSE®", a food grade crystalline trehalose hydrate, which was equal to about nine percent of sucrose with respect to the trehalose content, d.s.b., with and in 100 parts by weight of "COUPLING SUGAR®", a liquid saccharide composition, containing erlose and sucrose and having about 25% moisture and about 14% sucrose, d.s.b., commercialized by Hayashibara Shoji, Inc., Okayama, Japan. Comparing with the material liquid saccharide composition, the product thus obtained succeeded in improving the aftertaste inherent to sucrose in the product without altering the sweetening power of sucrose. Therefore, the product can be arbitrarily used to sweeten food products and used as a relatively-low dental-caries-inducing sweetener.

Example A-5

A liquid saccharide composition was obtained by mixing and dissolving 1.4 parts by weight of "TREHAOSE®", a food grade crystalline trehalose hydrate, which was equal to about 10% of the sucrose below with respect to the trehalose, d.s.b., with and in 100 parts by weight of "NYUKAOLIGO®", a liquid saccharide composition containing lactosucrose and sucrose and having about 25% moisture and about 17% sucrose, d.s.b., commercialized by Hayashibara Shoji, Inc., Okayama, Japan. Comparing with the material liquid saccharide composition, the product thus obtained succeeded in improving the aftertaste inherent to sucrose without lowering the sweetening power of sucrose. Therefore, the product can be arbitrarily used to sweeten food products and used as a saccharide for promoting the growth of bifid bacteria and for promoting the absorption of minerals in living bodies.

Example A-6

In 94 parts by weight of a 66 w/v % sucrose solution was dissolved by heating six parts by weight of "TREHAOSE®", a food grade crystalline trehalose hydrate, to obtain a liquid saccharide solution containing about 67 w/v % of sucrose and trehalose. The product is a stable and easily-handleable liquid saccharide and is free from solidification of sucrose and trehalose even at ambient temperature. The product can be arbitrarily used to sweeten food products because it has improved sharpness of aftertaste of sucrose without lowering the sweetening power of sucrose.

Example B-1Sweetener

One part by weight of a saccharide composition, obtained by the method in Example A-1, was mixed to homogeneity with 0.01 part by weight of "αG SWEET", α-glycosyl stevioside commercialized by Toyo Sugar Refining Co., Ltd., Tokyo, Japan, and 0.01 part by weight of "ASPARTAME", L-aspartyl-L-phenylalanine methyl ester, and the mixture was subjected to a granulator to obtain a granulated sweetener. The product has satisfactory quality of sweetness and favorable sharpness of aftertaste, as well as having three times higher sweetening power of sucrose and about one third lower calorific value of sucrose per sweetening power. Thus, the product can be favorably used as a relatively-low calorific sweetener for sweetening low-calorific food products for fat persons and diabetics who are restricted not to taking excessive calories.

Example B-2Canned coffee

One hundred parts by weight of roasted coffee beans was pulverized and extracted with about 1,000 parts by weight of heated water to obtain an about 860 parts by weight of an extract. An about 450 parts by weight of the extract was mixed to homogeneity with 80 parts by weight of a saccharide composition, obtained by the method in Example A-2, and about 550 parts by weight of water containing an adequate amount of sodium bicarbonate to obtain a coffee with a pH of about 7. The coffee was in a conventional manner injected into cans and sterilized by heating at 120°C for 30 min to obtain a canned coffee. The product has satisfactory flavor and taste, adequate sweetness, and favorable sharpness of aftertaste. Even if tasted after cooling in summer season, the product satisfactorily retains the flavor, taste, and high-quality.

Example B-3Chewing gum

5 Three parts by weight of gum base was heated and melted by heating up to be softened, admixed with five parts by weight of a saccharide composition, obtained by the method in Example A-3, and two parts by weight of crystalline maltitol anhydride, then further mixed with adequate amounts of a flavor and coloring agents. The mixture thus obtained was kneaded by a roll, shaped, and packaged into the desired product. The product is a chewing gum having satisfactory texture, flavor, and taste.

Example B-4Chocolate

15 Forty parts by weight of cacao paste, 10 parts by weight of cacao butter, 45 parts by weight of granulated sugar or sucrose, and five parts by weight of "TREHAOSE®", a food grade crystalline trehalose hydrate, which was equal to about 10% of the sucrose, d.s.b., were passed through a refiner to lower the granule size of the contents, placed in a conche, and kneaded up at 50°C for two days and nights. During the kneading step, 0.05 part by weight of lecithin was added to the contents and sufficiently kneaded and dispersed. The resulting product was set to 31°C by a ther-
 20 moregulator, poured into a mold just before solidifying the butter, deaerated by a vibrator, and passed through a 10°C cooled tunnel over about 20 min to solidify the contents. The solidified contents were removed from the mold and packaged into the desired product which has substantially no hygroscopicity but has a satisfactory color, gloss, and internal tissue, and mildly melts in your mouth to give favorable sweetness, flavor, taste, and sharpness of aftertaste.

Example B-5Sweetened condensed milk

30 In 100 parts by weight of fresh milk were dissolved one part by weight of a saccharide composition obtained by the method in Example A-1, and three parts by weight of a liquid saccharide composition obtained by the method in Example A-5, and the solution was sterilized by heating on a plate heater, concentrated up to give a concentration of 70%, and sterilely canned. The product has high-quality sweetness, flavor, taste, and sharpness of aftertaste, and it can be used to sweeten foods for infants, fruits, coffees, cocoas, and teas.

Example B-6An (a bean jam)

40 Ten parts by weight of material red beans was in a conventional manner boiled after mixed with water, followed by removing astringency, harshness, and water-soluble impurities to obtain an about 21 parts by weight of "azuki-tsubu-an" (a red bean jam with unpasted red beans). The an was mixed with 14 parts by weight of white superior soft sugar, 11 parts by weight of "TREHAOSE®", a food grade crystalline trehalose hydrate, which was equal to about seven percent of the sucrose, d.s.b., and four parts by weight of water. The mixture was heated, mixed with a small amount of salad oil, and kneaded up into an about 35 parts by weight of the desired product with unpasted red beans.
 45 The product has substantially no fading of color but has pleasure to the palate, as well as high-quality sweetness, flavor, taste, and sharpness of aftertaste. Thus, it can be suitably used as a material for bean-jam-buns, manju (a Japanese confectionery), dumplings, monaka (a bean-jam-filled wafer), ice creams, and sherbets.

Example B-7Custard cream

50 One hundred parts by weight of corn starch, 100 parts by weight of a saccharide composition obtained by the method in Example A-6, 100 parts by weight of maltose, and one part by weight of salt were sufficiently mixed, and the mixtur was admixed with 280 parts by weight of fresh eggs, followed by stirring. To the resulting mixture was gradually added 1,000 parts by weight of boiling milk and continued stirring over fire, then the heating was stopped
 55 when the whole contents completely gelatinized and became semi-transparency, followed by cooling the mixture, adding thereto an adequate amount of a vanilla flavor, then weighing, injecting and packing the contents to obtain the

desired product. The product has smooth gloss, mild sweetness and taste, and the retrogradation is satisfactorily inhibited, resulting in a long shelf-life by a large margin.

As evident from the above, the present invention improves the aftertaste of sucrose without lowering the sweetening power of sucrose by incorporating trehalose into sucrose in an amount of 2-12% to the sucrose, d.s.b. The present saccharide compositions and food products improve the aftertaste of sucrose without lowering the sweetening power. The present invention improves the aftertaste of sucrose now used as a sweetener in large quantities, and this would greatly influence, particularly, on the fields of food products such as sweetener, beverage, and processed-food industries, and it would be a great industrial significance.

While there has been described what is at present considered to be the preferred embodiments of the invention, it will be understood the various modifications may be made therein, and it is intended to cover in the appended claims all such modifications as fall within the true spirit and scope of the invention.

Claims

1. A method for improving the aftertaste of sucrose, which comprises a step of incorporating trehalose into sucrose in an amount of 2-12 w/w % to said sucrose, on a dry solid basis.
2. The method of claim 1, wherein said sucrose is a powdered sucrose and is prevented from solidification.
3. The method of claim 1 or claim 2, which is directed to a food product containing at least 10 w/w % sucrose, on a dry solid basis.
4. A saccharide composition, which contains trehalose in an amount of 2-12 w/w % to sucrose, on a dry solid basis.
5. The composition of claim 4, wherein the aftertaste of sucrose is improved without lowering the sweetening power of sucrose.
6. The composition of claim 4 or claim 5, wherein said sucrose is a member selected from the group consisting of white sugars, granulated sugars, white superior soft sugars, raw sugars, and liquid sugars.
7. The composition of any one of claims 4, 5 or 6, wherein said sucrose is a powdered sucrose and is prevented from solidification.
8. The composition of any one of claims 4 to 7, which contains at least 10 w/w % sucrose, on a dry solid basis, and optionally other saccharide excluding said trehalose.
9. The composition of claim 8, wherein said other saccharide is a member selected from the group consisting of monosaccharides, disaccharides excluding sucrose, trisaccharides, tetrasaccharides, sugar alcohols, and mixtures thereof.
10. The composition of claim 9, wherein said tri- and tetra-saccharides are oligosaccharides having a sucrose structure.
11. A method for producing food products, which comprises a step of incorporating the saccharide composition of claim 4 into a material for food products.
12. The method of claim 11, wherein at least 30 w/w % of the saccharide composition, on a dry solid basis, is incorporated into the material.
13. A food product wherein the aftertaste of sucrose is improved by incorporating the saccharide composition of claim 4 into said food product without lowering the sweetening powder of sucrose.
14. The food product of claim 13, which is a member selected from the group consisting of sweeteners, coffees, teas, chewing gums, chocolates, sweetened condensed milks, and bean jams.
15. The food product of claim 13 or claim 14, which contains at least 30 w/w % of the saccharide composition, on a dry solid basis.

16. A food product wherein the aftertaste of sucrose is improved without lowering the sweetening powder of sucrose by incorporating trehalose into said sucrose in an amount of 2-12 w/w % to said sucrose, on a dry solid basis.
17. The food product of claim 16, which is a member selected from the group consisting of sweeteners, coffees, teas, chewing gums, chocolates, sweetened condensed milks, and bean jams.
18. An agent for improving the aftertaste of sucrose, which comprises trehalose as an effective ingredient.
19. The agent of claim 18, which is for compositions containing sucrose.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 30 4315

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	EP 0 622 025 A (AJINOMOTO KK) 2 November 1994	1-19	A23L1/236
Y	* example 2 *	1-19	
X	DATABASE WPI Section Ch, Week 9737 Derwent Publications Ltd., London, GB; Class D13, AN 97-396989 XP002078090 & JP 09 173 017 A (LOTTE CO LTD) * abstract *	1-19	
Y	PATENT ABSTRACTS OF JAPAN vol. 097, no. 002, 28 February 1997 & JP 08 256694 A (NIPPON SHOKUHI KAKO CO LTD), 8 October 1996, * abstract *	1-19	
Y	PATENT ABSTRACTS OF JAPAN vol. 097, no. 003, 31 March 1997 & JP 08 298932 A (NIPPON SHOKUHI KAKO CO LTD), 19 November 1996, * abstract *	1-19	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int.Cl.6) A23L
Place of search MUNICH		Date of completion of the search 18 September 1998	Examiner Bendl, E
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03 92 (P4/C21)